

Vortices as degenerate metrics

J. M. Baptista

December 2012

Abstract

We note that the Bogomolny equation for abelian vortices is precisely the condition for invariance of the Hermitian-Einstein equation under a degenerate conformal transformation. This leads to a natural interpretation of vortices as degenerate hermitian metrics that satisfy a certain curvature equation. Using this viewpoint, we rephrase standard results about vortices and make some new observations. We note the existence of a conceptually simple, non-linear rule for superposing vortex solutions, and we describe the natural behaviour of the L^2 -metric on the moduli space upon certain restrictions.

Keywords: Vortex equation; degenerate Hermitian-Einstein metric; vortex superposition;

1 Vortices on Riemann surfaces

It is well known that solving the abelian vortex equation is equivalent to solving the so-called Kazdan-Warner equation [Bra]. The latter is a classical PDE that originally appeared in the problem of finding Riemannian metrics with prescribed scalar curvature [KW]. So in some sense abelian vortices should be related to Riemannian metrics. Our aim here is to find a natural way to formulate this relation. We first deal with the two-dimensional case of vortices living on a Riemann surface. In Section 2 we extend the discussion to vortices on higher-dimensional Hermitian manifolds and describe the relation between the vortex equation and the Hermitian-Einstein equation. Although we work with compact manifolds, a large part of the discussion is local, so follows through to the non-compact case.

The curvature equation

Let M be a compact Riemann surface equipped with a Kähler form ω . Let $L \rightarrow M$ be a hermitian line bundle over that surface and suppose that we are given a connection A and a non-trivial section ϕ of L such that

$$\bar{\partial}_A \phi = 0 . \quad (1)$$

This condition means that ϕ is a holomorphic section with respect to the holomorphic structure on L induced by the connection A . In particular, ϕ vanishes at isolated points $q_j \in M$ with a well defined, positive multiplicity $n_j \in \mathbb{Z}$. So a solution of (1) determines an effective divisor $D = \sum n_j q_j$ on the surface M . Consider now the degenerate Kähler form on the surface defined by

$$\omega' := \frac{1}{\tau} |\phi|^2 \omega , \quad (2)$$

where τ is a positive constant. This is a smooth 2-form on M that vanishes around each point q_j as $|z|^{2n_j}$, where z is a complex coordinate on M centered at q_j . We will say that a Kähler metric with this property is degenerate along the divisor D . Our basic question is: suppose that the pair (A, ϕ) satisfies the vortex equation

$$iF_A + e^2 (|\phi|^2 - \tau) \omega = 0 , \quad (3)$$

where e^2 is another positive constant and F_A denotes the curvature of the connection A . What does this imply for ω' ?

Proposition 1.1. *The pair (A, ϕ) satisfies the vortex equation (3) on M if and only if the metric ω' is degenerate along D and satisfies*

$$iF_{\omega'} + e^2 \tau \omega' = iF_{\omega} + e^2 \tau \omega , \quad (4)$$

over $M \setminus \{q_j\}$, where F_{ω} denotes the curvature form of the Riemannian metric ω .

Proof. The basic observation is that, away from the zeroes of ϕ , the curvatures are related by $F_A = \bar{\partial}\partial \log |\phi|^2 = F_{\omega'} - F_{\omega}$. See the proof of Proposition 2.1 for more details. $\square$

So the vortex equation is precisely the condition for the conformal transformation (2) to preserve the form $iF_{\omega} + e^2\tau\omega$. Studying abelian vortices on (M, ω) is the same thing as studying Kähler metrics ω' that satisfy equation (4). Abelian vortices define such degenerate metrics and, conversely, all vortex solutions are given by quotients of the form ω'/ω . The well known fact that there exists only one vortex solution (A, ϕ) with nowhere vanishing ϕ , up to gauge transformations, and that this solution has constant norm $|\phi|^2 = \tau$, just means that $\omega' = \omega$ is the unique non-degenerate solution to the curvature equation (4). Given an effective divisor $D = \sum n_j q_j$ on M , the well known fact that there exists only one vortex solution (A, ϕ) associated to D , up to gauge transformations [Bra, Gar, JT], tells us that there exists a unique metric ω' on the Riemann surface that satisfies (4) and is degenerate at the points $q_j \in M$ with multiplicity n_j .

Remark. Suppose that the metric ω on the surface has constant scalar curvature equal to $-e^2\tau$. Equation (4) then reduces to $iF_{\omega'} + e^2\tau\omega' = 0$. This means that each vortex solution on (M, ω) defines through (2) a degenerate hyperbolic metric ω' with the same curvature $-e^2\tau$. Conversely, all vortex solutions on (M, ω) can be obtained as quotients $\tau\omega'/\omega$ of hyperbolic metrics. This is a familiar fact when $M = \mathbb{H}$ is the hyperbolic half-plane. In this case, all degenerate hyperbolic metrics on $\mathbb{H}$ can be obtained as pullbacks $\omega' = f^*\omega$ of the Poincaré metric by Blaschke products $f : \mathbb{H} \rightarrow \mathbb{H}$. So all vortex solutions can be explicitly written as quotients $(f^*\omega)/\omega$, a fact that was originally noted by Witten [Wi]. When (M, ω) is a compact hyperbolic surface, Manton and Rink noted that one can also obtain vortex solutions as quotients $(f^*\omega_Y)/\omega$, where (Y, ω_Y) is a second hyperbolic surface and $f : M \rightarrow Y$ is a ramified holomorphic map [MR]. Unfortunately these maps are rare, so only isolated and non-explicit vortex solutions can be obtained in this way (an upset that improves marginally if we accept vortices with singularities [BB]). Our observation here is that, even in the compact case, all vortex solutions can still be written as quotients of hyperbolic metrics on M . The difficulty is that the degenerate metrics $f^*\omega_Y$ provided by ramified maps represent only a tiny subset of all the degenerate hyperbolic metrics on M .

Degenerate metrics

A word now about standard properties of degenerate metrics. Firstly, observe that the Riemannian curvature form $F_{\omega'}$ has a smooth extension to the degenerate points, even though the local 1-forms $A_{\omega'}$ of the Chern connection do not. In fact, around each $q_j \in M$ we can write

$$\omega' = |z|^{2n_j} f_j \omega, \quad (5)$$

where f_j is a smooth and positive function locally defined around q_j . It follows that the Levi-Civita (or Chern) connection is given by

$$A_{\omega'} = A_{\omega} + n_j z^{-1} dz + \partial(\log f_j)$$

in a local holomorphic trivialization of TM . So the connection is singular at the degenerate points and the residues n_j can be probed by integrating $A_{\omega'}$ around small circles centered at the q_j 's. On the other hand, the curvature

$$F_{\omega'} = dA_{\omega'} = F_{\omega} + \bar{\partial}\partial(\log f_j)$$

can be smoothly extended to the degenerate points q_j . Note that the last term in the equation above, although locally exact, is not globally exact, since the functions f_j are not globally defined. In fact, one can check that if ω' is defined by (2) and the pair (A, ϕ) solves the holomorphy equation (1), then away from the zeroes of ϕ we have

$$F_{\omega'} = F_{\omega} + F_A, \quad (6)$$

where, again, F_A is the curvature of the $U(1)$ -connection A . This identity is another way to recognize that $F_{\omega'}$ can be smoothly extended to the whole M . However, note that this smooth extension is not globally a curvature form, because the Levi-Civita connection has singularities. In particular, one cannot directly apply Stokes' theorem to $F_{\omega'}$ over domains that contain degenerate points. The Gauss-Bonnet theorem in its usual form is also not applicable to the smooth extension of $F_{\omega'}$, as the integral of identity (6) yields

$$\frac{i}{2\pi} \int_{M \setminus \{q_j\}} F_{\omega'} = (2 - 2g) + \sum_j n_j, \quad (7)$$

where g is the genus of M . To remedy these undesirable features, one usually considers an extension of the curvature $F_{\omega'}$ to the degenerate points that is different from the smooth one. This second extension regards the curvature of the degenerate metric as an integration current, and is defined by

$$F_{\omega'} := \begin{cases} dA_{\omega'} & \text{over } M \setminus \{q_j\} \\ 2\pi i n_j \delta(q_j) & \text{at each point } q_j, \end{cases} \quad (8)$$

where $\delta(q_j)$ is the delta function. Thus, at the price of considering distributions, the usual formulae are applicable to the curvature, and we have for instance

$$\frac{i}{2\pi} \int_M F_{\omega'} = (2 - 2g).$$

Defining the curvature of a degenerate metric to have delta functions at the points q_j also makes sense if we think of the degenerate metric as a limit of non-degenerate metrics. In this case one can check that the curvature of the non-degenerate metrics increases at

q_j as these metrics tend to zero at q_j , and the limit will be a delta function. Moreover, since the curvature of all the non-degenerate metrics satisfies Gauss-Bonnet, it is natural to define a limit curvature $F_{\omega'}$ that also satisfies Gauss-Bonnet. Observe that if we use integration currents on M , the curvature equation (4) should be rewritten as

$$iF_{\omega'} + e^2\tau\omega' = iF_{\omega} + e^2\tau\omega - 2\pi \sum_j n_j \delta(q_j) , \quad (9)$$

and this is an equation over the whole surface. Comparing with (4), we see that the delta functions encode the boundary conditions for ω' at the degenerate points. They play exactly the same role here as in the original Taubes' equation for abelian vortices [JT].

Minimal volume bound

Integrating the curvature equation (4) over $M \setminus \{q_j\}$, or integrating equation (9) over M , we see that each solution ω' must satisfy

$$\text{Vol}(M, \omega') = \text{Vol}(M, \omega) - \frac{2\pi}{e^2\tau} \sum_j n_j .$$

Thus each additional degeneracy (i.e. vortex) reduces the volume of the metric ω' by $2\pi/(e^2\tau)$. Since any Riemannian metric with a finite number of degenerate points certainly has positive volume, we also conclude that a necessary condition for the existence of solutions of the curvature equation is that

$$2\pi \sum_j n_j < e^2\tau \text{Vol}(M, \omega) .$$

This, of course, is just the standard condition for the existence of vortex solutions associated to a non-zero divisor on M [Bra].

Remark. The interpretation of vortices as a deformation of the background Riemannian metric satisfies basic intuitive properties. For instance, it is known that the rescaling factor $\tau^{-1}|\phi|^2$ tends exponentially fast to the vacuum value 1 as one gets away from the vortex core, at least for vortices on the flat plane [JT]. So the background metric ω is substantially deformed only in a very localized region around the vortices. Moreover, an application of the maximum principle says that the conformal factor is bounded by $0 \leq \tau^{-1}|\phi|^2 \leq 1$ over the entire surface M [JT, Bra]. Hence the total deformation of the background metric ω can be measured by the difference of volumes $\text{Vol}(M, \omega) - \text{Vol}(M, \omega')$, which is proportional to the vortex number.

Hidden symmetries and vortex superposition

The interpretation of abelian vortices as degenerate metrics exposes two symmetries of the vortex equations that are less manifest in their usual form. The first is the fact that

the curvature equation (4) is essentially symmetric in ω and ω' . If ω' is a solution on the background (M, ω) , then ω is a solution on the background (M, ω') . To interpret this in terms of the usual vortex equations requires working with meromorphic sections ϕ . The observation is that if (L, ϕ, A) is a solution of (3) on (M, ω) , then $(L^{-1}, \phi^{-1}, -A)$ is a solution on the background $(M, \tau^{-1}|\phi|^2\omega)$. Meromorphic sections appear because the boundary conditions at the degenerate points make the curvature equation not exactly symmetric in ω' and ω . This is most manifest in the form (9) of the equation, where an interchange of the two metrics requires an accompanying change of sign of the divisor in order to preserve the equation.

The second hidden symmetry is a transitivity property. If ω' is a solution of the curvature equation (4) on the background (M, ω) and ω'' is a solution on the background (M, ω') , then ω'' is a solution on the background (M, ω) . This is evident from

$$iF_{\omega''} + e^2\tau\omega'' = iF_{\omega'} + e^2\tau\omega' = iF_{\omega} + e^2\tau\omega.$$

In the usual formulation of vortices, this transitivity corresponds to the observation that if (L_1, ϕ_1, A_1) is a vortex solution on (M, ω) and (L_2, ϕ_2, A_2) is a vortex solution on the degenerate background $(M, \tau^{-1}|\phi_1|^2\omega)$, then the tensor product of hermitian bundles $(L_2 \otimes L_1, \tau^{-1/2}\phi_2\phi_1, A_2 + A_1)$ is a vortex solution on the original background (M, ω) . It is straightforward to check this directly using equation (3). The basic identity is

$$iF_{A_2+A_1} + e^2(\tau^{-1}|\phi_1\phi_2|^2 - \tau)\omega = iF_{A_2} + e^2(|\phi_2|^2 - \tau)(\tau^{-1}|\phi_1|^2\omega) + [iF_{A_1} + e^2(|\phi_1|^2 - \tau)\omega].$$

This property of vortex solutions is very appealing and, apparently, has not been remarked before. It says that we can construct a (d_1+d_2) -vortex solution on (M, ω) by first obtaining a d_1 -vortex solution on (M, ω) , then deforming the background ω to the degenerate metric $\tau^{-1}|\phi_1|^2\omega$ dictated by that first solution, and finally solving the equation for d_2 -vortices on the deformed background. We can also do it the other way around, of course, solving for d_1 -vortices on the background deformed by d_2 -vortices. This amounts to a conceptually simple, non-linear rule for superposing vortices. Again, it states that instead of naively multiplying vortex solutions obtained on the same background, the correct prescription to superpose vortices implies using a second solution obtained on the background deformed by the first solution. If one knew how to solve the 1-vortex equation on any background, a recursive application of this rule would yield the d -vortex solution on any background.

Energy functional

Another basic fact about the vortex equations is that they can be derived as the (Bogomolny) condition for static, minimal energy configurations of the abelian Higgs model at critical coupling [MS, Bra]. So they minimize the energy functional

$$E(A, \phi) = \int_{(M, \omega)} \frac{1}{2e^2} |F_A|^2 + |d_A\phi|^2 + \frac{e^2}{2} (|\phi|^2 - \tau)^2.$$

One may wonder how to rephrase this in terms of degenerate metrics. Given an effective divisor $D = \sum n_j q_j$ on a surface (M, ω) , we define the energy density of a Kähler metric ω' that degenerates along D by

$$\mathcal{E}(\omega') := \frac{1}{2e^2} |F_{\omega'} - F_\omega|^2 + 2\tau |\mathrm{d}\sqrt{\omega'/\omega}|^2 + \frac{e^2\tau^2}{2} |\omega' - \omega|^2 \quad (10)$$

This is a smooth function on $M \setminus \{q_j\}$. It can be rewritten as

$$\mathcal{E}(\omega') = \frac{1}{2e^2} |F_{\omega'} - F_\omega - ie^2\tau(\omega' - \omega)|^2 + i\tau * [F_{\omega'} - F_\omega + \partial\bar{\partial}(\omega'/\omega)]$$

over that domain, where $*$ is the Hodge operator with respect to ω . Using equation (7) and Stokes' theorem, it is clear that the total energy functional satisfies

$$E(\omega') := \int_{M \setminus \{q_j\}} \mathcal{E}(\omega') \omega \geq 2\pi\tau \sum_j n_j,$$

with the equality standing precisely when ω' satisfies the curvature equation (4) over $M \setminus \{q_j\}$. This is the analog of the usual Bogomolny argument for vortices.

Remark. At first sight one could be worried that the square root term appearing in the energy density (10) may lead to singularities at the points where ω' vanishes. However, a direct calculation shows that if ω' vanishes at q_j as $|z|^{2n_j}$, the squared norm $|\mathrm{d}\sqrt{\omega'/\omega}|^2$ is smooth at that point for any integer $n_j \geq 0$.

Remark. In this discussion we have fixed the degeneration divisor D , have only considered metrics ω' with the boundary conditions prescribed by D , and the integrals are over the domain $M \setminus \mathrm{supp} D$. An equivalent approach would be to consider the curvature $F_{\omega'}$ as an integration current over the entire M , as in (8), and then encode the boundary conditions in delta functions appearing in the energy density. In this case one should substitute the term $F_{\omega'} - F_\omega$ in the formulae above by $F_{\omega'} - F_\omega + 2\pi\delta(D)$ and perform the integrations over the whole M .

Metric on the moduli space

It is well known that for each effective divisor $D = \sum n_j q_j$ on the surface M there is exactly one solution of the vortex equations (1) and (3), up to $U(1)$ -gauge transformations, such that D is the zero-set divisor of the Higgs field ϕ . So if we fix the degree of D to be d and let the points q_j vary in M , we get a moduli space $\mathcal{M}^d$ of vortex solutions that is isomorphic to the symmetric product $\mathrm{Sym}^d M$ of the surface. In terms of degenerate metrics, $\mathcal{M}^d$ is the space of solutions of the curvature equation (4) such that the Kähler metric ω' has d degenerate points on M , counting multiplicities.

The space $\mathcal{M}^d$ has a natural Kähler metric induced by its interpretation as a vortex moduli space [MS]. Using the language of [Ba, Sect.7], the corresponding Kähler form

$\omega_{\mathcal{M}}$ can be written in terms of the degenerate metrics as

$$\omega_{\mathcal{M}} = \frac{-1}{2e^2} \left\{ \int_M \partial_z \left[(\partial_{\bar{u}^k} \log f) (\partial_{\bar{z}} \partial_{u^j} \log f) \right] dz \wedge d\bar{z} \right\} du^j \wedge d\bar{u}^k, \quad (11)$$

where $f(\omega', p) := (\omega'/\omega)(p)$ is a non-negative smooth function on the cartesian product $\mathcal{M}^d \times M$, and we chose arbitrary local complex coordinates z on M and $\{u^k\}$ on $\mathcal{M}^d$. Observe that in (11) the integrand 2-form is exact outside the points where $\log f$ explodes, i.e. outside the degenerate points of the metric ω' . By Stokes' theorem, the integral therefore localizes to a sum of integrals around small circles centered at these degenerate points, which of course is just the standard Samols' localization for vortices [Sa, Ba].

Formula (11) and the superposition rule for vortices imply that the metric $\omega_{\mathcal{M}}$ on the moduli space has an interesting property upon restrictions, which we now explain. Let $\mathcal{M}_p^d \subset \mathcal{M}^d$ be the complex submanifold determined by fixing one vortex at position $p \in M$, i.e. $\mathcal{M}_p^d$ is the subset of unordered multiples in $\text{Sym}^d M$ of the form $(x_1, \dots, x_{d-1}, p)$ with varying $x_k \in M$. As a complex manifold, $\mathcal{M}_p^d$ clearly is isomorphic to $\mathcal{M}^{d-1}$. However, the Kähler metric on $\mathcal{M}_p^d$ induced by the embedding in $(\mathcal{M}^d, \omega_{\mathcal{M}})$ is different from the natural metric on $\mathcal{M}^{d-1}$ regarded as the moduli space of $d-1$ vortices. In other words, the remaining $d-1$ vortices “feel” the presence of the vortex fixed at p . But how, precisely, does this presence translate into the metric on $\mathcal{M}_p^d$? In view of superposition rule discussed before, a natural guess is that the vortex fixed at p deforms the background metric ω on the surface M to another metric ω'' , a metric that is degenerate at p , and then the remaining $d-1$ vortices behave as free vortices on the surface (M, ω'') . To be more precise:

Proposition 1.2. *Let ω'' be the unique solution of the curvature equation (4) on (M, ω) with a single simple degeneracy at the point $p \in M$. Then the submanifold $\mathcal{M}_p^d$ equipped with the metric inherited from $(\mathcal{M}^d, \omega_{\mathcal{M}})$ is isometric to the moduli space $\mathcal{M}^{d-1}$ of $d-1$ vortices living on the background surface (M, ω'') .*

To show that this is true, take a local chart $\{u^k\}$ of the moduli space $\mathcal{M}^d$ adapted to the submanifold $\mathcal{M}_p^d$. This means that in this chart $\mathcal{M}_p^d$ is determined by the equation $u^d = 0$ and $\{u^1, \dots, u^{d-1}\}$ are good complex coordinates on $\mathcal{M}_p^d$. Now, for any solution $\omega' \in \mathcal{M}_p^d$ of the curvature equation on (M, ω) , we can always factorize

$$\frac{\omega'}{\omega} = \frac{\omega'}{\omega''} \frac{\omega''}{\omega}.$$

Crucially, the factor ω''/ω is constant on the submanifold $\mathcal{M}_p^d$, it does not depend on the positions of the remaining $d-1$ vortices. So for the coordinates $\{u^1, \dots, u^{d-1}\}$ we have

$$\partial_{u^j} \log(\omega'/\omega) = \partial_{u^j} \log(\omega'/\omega'')$$

over $\mathcal{M}_p^d$, and the same applies to the anti-holomorphic derivatives. Thus the restriction of the Kähler form (11) to the submanifold $\mathcal{M}_p^d$ is given by the same formula (11), except

that we only use the coordinates $\{u^1, \dots, u^{d-1}\}$ and the function f is now ω'/ω'' . This coincides with the formula for the Kähler form on the moduli space $\mathcal{M}^{d-1}$ of $d-1$ vortices living on the background surface (M, ω'') .

There is an obvious extension of this property to the lower dimensional submanifolds $\mathcal{M}_{p_1, \dots, p_l}^d$ of the moduli space $\mathcal{M}^d$ determined by fixing the positions of $l < d$ vortices. It says that $\mathcal{M}_{p_1, \dots, p_l}^d$ equipped with the metric inherited from $(\mathcal{M}^d, \omega_{\mathcal{M}})$ is isometric to the moduli space $\mathcal{M}^{d-l}$ of $d-l$ vortices living on the background surface (M, ω'') , where this time ω'' is the unique solution of the curvature equation on (M, ω) with simple degeneracies at the points $p_1, \dots, p_l$.

Vortices with parabolic singularities

So far we have only considered degenerate metrics ω' on M that vanish with a positive integral power $|z|^{2n_j}$ at the degenerate points q_j . However, pretty much the whole story can be extended to metrics that vanish as $|z|^{2\alpha_j}$ for real numbers $\alpha_j > 0$. These correspond to abelian vortices on (M, ω) with parabolic singularities.

Suppose that we are given a divisor $\sum \alpha_j q_j$ on M with positive real coefficients. Denote by $[\alpha_j] \in \mathbb{Z}$ the integral part of α_j , and call $L \rightarrow M$ the complex line bundle of degree $\sum_j [\alpha_j]$. Let h be a hermitian metric on L with parabolic singularities of weight $\alpha_j - [\alpha_j]$ at the points q_j . This means that if s is a smooth local section of L that does not vanish at q_j , the squared norm of s is of the form

$$|s|_h^2 = |z|^{2(\alpha_j - [\alpha_j])} f_j(z, \bar{z})$$

for some smooth and positive function f_j defined around q_j . So h is degenerate at the points q_j , and all unitary connections on (L, h) will have logarithmic singularities at those points. However, this degeneracy does not affect the existence of vortex solutions. There still exists a smooth section ϕ of L and a unitary connection A that solve the vortex equations (1) and (3) over $M \setminus \{q_j\}$, and such that ϕ vanishes precisely at points q_j with multiplicity $[\alpha_j]$. Moreover, this pair (A, ϕ) is unique up to $U(1)$ -gauge transformations on M [BiGa]. Observe that for this solution the squared norm $|\phi|_h^2$ vanishes as $|z|^{2\alpha_j}$ at the point q_j . Therefore the Riemannian metric $\omega' := \tau^{-1} |\phi|_h^2 \omega$ satisfies the curvature equation (4) over $M \setminus \{q_j\}$ and is degenerate along the divisor $\sum \alpha_j q_j$, as required. All in all, we see that studying vortices on (M, ω) with parabolic singularities is the same thing as studying solutions ω' of (4) that vanish with positive real weights. All the formulae presented above remain valid if we substitute the integers n_j by the real weights α_j .

Remark. The existence and uniqueness of vortex solutions, and hence of solutions to (4) with prescribed degeneracies, has also been extended to the case where the background metric ω is allowed to have degeneracies or conical singularities, meaning that at a finite number of points $p_k \in M$, possibly coincident with the q_j 's, the background Kähler form ω behaves as $|z|^{2\beta_k}$ with real weights $\beta_k > -1$ [BB].

2 Vortices in higher dimensions

In the previous Section we have seen how the vortex equation (3) can be regarded as the condition for the conformal transformation (2) to preserve the form $iF_\omega + e^2\tau\omega$. Observe that the metric ω plays here a double role, both as a Riemannian metric on M and as a hermitian metric on TM with curvature F_ω . If we separate these roles and substitute TM by an arbitrary holomorphic vector bundle, it becomes clear that the vortex equation is in fact the condition for invariance of the general Hermitian-Einstein equation under a conformal transformation. Moreover, this is true on any Hermitian manifold M , not just on Riemann surfaces.

Let (M, ω) be a compact Hermitian manifold and let $L \rightarrow M$ be a hermitian line bundle. Suppose that we are given a connection A and a non-trivial section ϕ of L such that

$$\bar{\partial}_A \phi = 0, \quad F_A^{0,2} = 0. \quad (12)$$

Then ϕ is a holomorphic section with respect to the holomorphic structure on L induced by the connection A . In particular, its zero set determines an effective divisor $D = \sum_j n_j Z_j$ of analytic hypersurfaces on M . The squared norm $|\phi|^2$ is a smooth function on M that vanishes along the support of D . Now let $V \rightarrow M$ be a holomorphic vector bundle equipped with a hermitian metric f . We can consider the double rescaling

$$f' := \frac{1}{\tau'} |\phi|^{2\alpha} f \quad \text{and} \quad \omega' := \frac{1}{\tau} |\phi|^2 \omega \quad (13)$$

for any exponent $\alpha \neq 0$. Then ω' , for instance, is a smooth 2-form on M that vanishes at each hypersurface Z_j as $|z|^{2n_j}$, where z is a complex coordinate on M such that Z_j is locally defined by the equation $z = 0$. In the language of Section 1, the degeneracies of ω' and f' are prescribed by the divisors D and αD , respectively. Once again, our question is: suppose that the pair (A, ϕ) satisfies the vortex equation

$$i\Lambda_\omega F_A + e^2 (|\phi|^2 - \tau) = 0, \quad (14)$$

where $\Lambda_\omega F_A$ denotes the contraction of the curvature of A with the Kähler form ω on M . What does this imply for the metrics f' and ω' ?

Proposition 2.1. *The pair (A, ϕ) satisfies the vortex equation (14) on M if and only if the metrics ω' and f' defined by the conformal transformation (13) satisfy the equation*

$$(i\Lambda_{\omega'} F_{f'} + e^2 \alpha \tau) \omega' = (i\Lambda_\omega F_f + e^2 \alpha \tau) \omega \quad (15)$$

over $M \setminus \cup_j Z_j$, where F_f denotes the curvature of the Chern connection on (V, f) .

In other words, the abelian vortex equation (14) is precisely the condition for invariance of the form $(i\Lambda_\omega F_f + e^2 \alpha \tau) \omega$ under the conformal rescaling (13). The existence of smooth vortex solutions has been proved by Bradlow in the case of Kähler (M, ω) [Bra], and by Lupascu in the case of Hermitian manifolds [Lu].

Remark. Choose $V = TM$, $f = \omega$ and $\alpha = 1$. Observe that if the initial metric ω satisfies the Hermitian-Einstein condition $i\Lambda_\omega F_\omega + e^2\tau I = 0$, then each vortex solution on (M, ω) gives us a new degenerate metric ω' on M that satisfies the same condition. Taking the trace in equation (15), we see that if the initial metric ω has constant scalar curvature $s_\omega = \text{Tr}(i\Lambda_\omega F_\omega) = -e^2\tau \dim_{\mathbb{C}} M$ (not necessarily being Hermitian-Einstein), then each vortex solution on (M, ω) gives us a new degenerate metric ω' with the same scalar curvature. However, note that if (M, ω) is a higher-dimensional Kähler manifold, the rescaled forms ω' are no longer d-closed for non-trivial ϕ , and hence will only define Hermitian metrics on M .

Proof of Proposition 2.1. Let h denote the background hermitian metric on L . Since the pair (A, ϕ) satisfies equations (12), there exists a holomorphic structure on L such that A coincides with the Chern connection and ϕ is a holomorphic section. In a local holomorphic trivialization of L , the curvature of the Chern connection A is given by $F_A = \bar{\partial}\partial(\log h)$. Since ϕ is holomorphic, away from its vanishing set we have that $\bar{\partial}\partial \log(\phi\bar{\phi}) = 0$. Hence on $M \setminus \cup_j Z_j$ the curvature is given by $F_A = \bar{\partial}\partial(\log |\phi|_h^2)$. On the other hand, in a local holomorphic trivialization of the vector bundle V , the curvature of the Chern connection is given by the analogous formula $F_f = \bar{\partial}[(\partial f)f^{-1}]$, where the hermitian metric f should be locally regarded as a square matrix of dimension equal to the rank of V [CCL, p. 256]. In particular

$$\begin{aligned} F_{f'} - F_f &= \bar{\partial}[(\partial f')(f')^{-1} - (\partial f)f^{-1}] = \bar{\partial}[(\partial f')f^{-1}(f'f^{-1})^{-1} + (f'f^{-1})^{-1}f'(\partial f^{-1})] \\ &= \bar{\partial}\{[(\partial f')f^{-1} + f'(\partial f^{-1})](f'f^{-1})^{-1}\} = \bar{\partial}\partial \log(f'f^{-1}) = \alpha F_A I, \end{aligned}$$

where I stands for the identity automorphism of V and we have used that $f'f^{-1} = |\phi|^{2\alpha}(\tau')^{-1}I$ commutes with all matrices. Thus on $M \setminus \cup_j Z_j$ we have that

$$\left[i\Lambda_\omega F_A + e^2(|\phi|^2 - \tau) \right] \alpha \omega = i(\Lambda_{\omega'} F_{f'}) \omega' - i(\Lambda_\omega F_f) \omega + e^2 \alpha \tau (\omega' - \omega),$$

and the result follows. $\square$

The curvature equation (15) has the same symmetry and transitivity properties that were discussed in the case of vortices on Riemann surfaces. It follows that the superposition rule of Section 1 is still valid for abelian vortices over higher-dimensional M . The relevant identity is in this case

$$\begin{aligned} \left[i\Lambda_\omega F_{A_2+A_1} + e^2(\tau^{-1}|\phi_1\phi_2|^2 - \tau) \right] \omega &= \left[i\Lambda_{\omega'} F_{A_2} + e^2(|\phi_2|^2 - \tau) \right] \omega' + \\ &\quad + \left[i\Lambda_\omega F_{A_1} + e^2(|\phi_1|^2 - \tau) \right] \omega, \end{aligned}$$

where ω' denotes the deformed hermitian metric $\tau^{-1}|\phi_1|^2\omega$.

Remark. Just as in the case of Riemann surfaces, if we regard the curvature $F_{f'}$ of a degenerate metric as an integration current, the domain of equation (15) can be extended

to the entire manifold M . In this case one should add a delta function term the curvature equation, a term that encodes the boundary conditions in (15). The equation over M then becomes

$$(i\Lambda_{\omega'} F_{f'} + e^2 \alpha \tau) \omega' = (i\Lambda_{\omega} F_f + e^2 \alpha \tau) \omega - 2\pi \delta_D, \quad (16)$$

where δ_D denotes the current of integration over the divisor D . That this is the correct term to add, is a consequence of the Poincaré-Lelong formula $\bar{\partial}\partial(\log|\phi|^2) = F_A + 2\pi i \delta_D$ applied to the line bundle L .

The modified vortex equation

As before, let $L \rightarrow M$ be a hermitian line bundle and let (A, ϕ) be a pair that satisfies the holomorphy equations (12). We assume that ϕ is non-trivial, so that its zero set determines an effective divisor $D = \sum_j n_j Z_j$ on M . A modified version of the vortex equation has been often considered, namely

$$i\Lambda_{\omega} F_A + \frac{1}{\dim_{\mathbb{C}} M} (|\phi|^2 + s_{\omega}) = 0, \quad (17)$$

where $s_{\omega} = i \operatorname{Tr}(\Lambda_{\omega} F_{\omega})$ denotes the scalar curvature on (M, ω) . When M is a Kähler surface, this equation is equivalent to a perturbation of the Seiberg-Witten equations [BG]. For compact Kähler (M, ω) , it can be shown that the existence of solutions to (17) depends only on the average value

$$\bar{s}_{\omega} := \frac{1}{\operatorname{Vol} M} \int_{(M, \omega)} s_{\omega}$$

of the scalar curvature of ω . Solutions (A, ϕ) to (17) exist if and only if the vortex equation (14) with $\tau = -\bar{s}_{\omega}$ has solutions [BG]. In particular, we must have $\bar{s}_{\omega} < 0$ for non-trivial solutions to exist. For a general compact Hermitian manifold (M, ω) , the existence condition is slightly more evolved [Lu]. Equation (17) also has a very simple interpretation in terms of degenerate metrics. As usual, for given data (L, M, ω, A, ϕ) , define the deformed metric

$$\omega' := \frac{-1}{\bar{s}_{\omega}} |\phi|^2 \omega.$$

Proposition 2.2. *The pair (A, ϕ) satisfies the modified vortex equation (17) on M if and only if the degenerate metric ω' has constant scalar curvature $s_{\omega'} = \bar{s}_{\omega}$ over $M \setminus \cup_j Z_j$.*

Proof. If $K \rightarrow M$ denotes the canonical line bundle, the induced hermitian metric on K rescales as

$$\det \omega' = \left(\frac{-1}{\bar{s}_{\omega}} |\phi|_h^2 \right)^{\dim_{\mathbb{C}} M} \det \omega.$$

Using that $F_A = \bar{\partial}\partial(\log|\phi|^2)$ away from the vanishing set of ϕ , as before, it is clear that the real Ricci form transforms as

$$\rho_{\omega'} = i\bar{\partial}\partial\log(\det\omega') = i(\dim_{\mathbb{C}} M) F_A + \rho_{\omega}$$

over $M \setminus \cup_j Z_j$. Thus over this domain we get that

$$\left[i\Lambda_{\omega} F_A + \frac{1}{\dim_{\mathbb{C}} M} (|\phi|_h^2 + s_{\omega}) \right] \omega = \frac{1}{\dim_{\mathbb{C}} M} \left[(\Lambda_{\omega'} \rho_{\omega'}) \omega' - (\Lambda_{\omega} \rho_{\omega}) \omega - \bar{s}_{\omega} \omega' + s_{\omega} \omega \right]$$

and the result follows. $\square$

Acknowledgements. It is a pleasure to thank Dennis Eriksson and Nick Manton for helpful discussions about integration currents and vortex superposition, respectively. I also thank CAMGSD-IST for a generous fellowship.

References

- [Ba] J. Baptista: On the L^2 -metric of vortex moduli spaces, *Nucl. Phys. B* **844** (2011), 308–333.
- [BB] J. Baptista and I. Biswas: Abelian vortices with singularities, arXiv:1207.0863.
- [BiGa] O. Biquard and O. García-Prada: Parabolic vortex equations and instantons of infinite energy, *Jour. Geom. Phys.* **21** (1997), 238–254.
- [Bra] S. Bradlow: Vortices in holomorphic line bundles over closed Kähler manifolds, *Commun. Math. Phys.* **135** (1990), 1–17.
- [BG] S. Bradlow and O. García-Prada: Non-abelian monopoles and vortices, *Geometry and Physics*, Lecture Notes in Pure and Appl. Math. **184**, 567–589, Dekker, 1997.
- [CCL] S. Chern, W. Chen and K. Lam: *Lectures on Differential Geometry*, World Scientific, 1998.
- [Gar] O. García-Prada: Invariant connections and vortices, *Commun. Math. Phys.* **156** (1993), 527–546.
- [JT] A. Jaffe and C. Taubes: *Vortices and monopoles*, Birkhauser, 1980.
- [KW] J. Kazdan and F. Warner: Curvature functions for compact 2-manifolds, *Annals of Math.* **99** (1974), 14–47.

- [Lu] P. Lupascu: The abelian vortex equations on Hermitian manifolds, *Math. Nachr.* **230** (2001), 99–115.
- [MR] N. Manton and N. Rink: Vortices on hyperbolic surfaces, *Jour. Phys. A* **43** (2010), 434024.
- [MS] N. Manton and P. Sutcliffe: *Topological Solitons*, Cambridge Univ. Press, 2004.
- [Sa] T. Samols : ‘*Vortex scattering*’; Commun. Math. Phys. **145** (1992), 149–179.
- [Wi] E. Witten: Some exact multipseudoparticle solutions of classical Yang-Mills theory, *Phys. Rev. Lett.* **38** (1977), 121–124.

CENTRE FOR MATHEMATICAL ANALYSIS, GEOMETRY, AND DYNAMICAL SYSTEMS (CAMGSD),
 INSTITUTO SUPERIOR TÉCNICO, AV. ROVISCO PAIS, 1049-001 LISBON, PORTUGAL

Email address: joao.o.baptista@gmail.com